

## NONEXPANSIVE MAPPINGS ON HILBERT'S METRIC SPACES

BAS LEMMENS

---

ABSTRACT. This paper deals with the iterative behavior of nonexpansive mappings on Hilbert's metric spaces  $(X, d_X)$ . We show that if  $(X, d_X)$  is strictly convex and does not contain a hyperbolic plane, then for each nonexpansive mapping, with a fixed point in  $X$ , all orbits converge to periodic orbits. In addition, we prove that if  $X$  is an open 2-simplex, then the optimal upper bound for the periods of periodic points of nonexpansive mappings on  $(X, d_X)$  is 6. The results have applications in the analysis of nonlinear mappings on cones, and extend work by Nussbaum and others.

### 1. Introduction

In [11] Hilbert introduced the following metric spaces which generalize the Cayley–Klein model of the hyperbolic plane. Let  $X \subseteq \mathbb{R}^n$  be a bounded open convex set. For  $x \neq y$  in  $X$  define the distance between  $x$  and  $y$  to be the logarithm of the *cross-ratio*,

$$[a, x, y, b] = \frac{|ay| |bx|}{|ax| |by|},$$

where  $a$  and  $b$  are the points of intersection of the straight-line through  $x$  and  $y$  and the (Euclidean) boundary,  $\partial X$ , of  $X$  such that  $x$  is between  $a$  and  $y$ , and  $y$  is between  $b$  and  $x$ , see Figure 1.

---

2010 *Mathematics Subject Classification*. Primary 47H09, 47H07; Secondary 54H20.

*Key words and phrases*. Nonexpansive mappings, Hilbert's metric, periodic orbits, hyperbolic plane, simplices, nonlinear mappings on cones.




FIGURE 1. Hilbert's metric

So, for  $x \neq y$  in  $X$ ,

$$d_X(x, y) = \log[a, x, y, b].$$

The metric  $d_X$  is called *Hilbert's metric* on  $X$ . In case  $X$  is the interior of an ellipse,  $(X, d_X)$  is a model for the hyperbolic plane.

Hilbert's metric is a natural example of a projective metric, meaning that straight-line segments are geodesics, and plays a role in the solution of Hilbert's fourth problem [1]. In this paper we study the iterates of *nonexpansive mappings*  $g: X \rightarrow X$  on Hilbert's metric spaces, so

$$d_X(g(x), g(y)) \leq d_X(x, y) \quad \text{for all } x, y \in X.$$

Motivating examples arise from nonlinear mappings on cones. More precisely, let  $C \subseteq \mathbb{R}^{n+1}$  be a closed cone with non-empty interior  $C^\circ$ , and let  $C^* = \{\phi \in \mathbb{R}^{n+1}: \langle x, \phi \rangle \geq 0 \text{ for all } x \in C\}$  denote the *dual cone*. The cone  $C$  induces a partial ordering  $\leq_C$  on  $\mathbb{R}^{n+1}$  by  $x \leq_C y$  if  $y - x \in C$ . Suppose that  $\phi$  is in the interior of  $C^*$  and let  $X_\phi = \{x \in C^\circ: \langle x, \phi \rangle = 1\}$ , which is a bounded, convex, relatively open set. Now if  $f: C^\circ \rightarrow C^\circ$  is a monotone mapping, i.e.  $f$  preserves  $\leq_C$  on  $C^\circ$ , and  $f$  is homogeneous of degree 1, then  $g: X_\phi \rightarrow X_\phi$  given by,

$$(1.1) \quad g(x) = \frac{f(x)}{\langle f(x), \phi \rangle} \quad \text{for all } x \in X_\phi,$$

is nonexpansive under Hilbert's metric, see [5], [22]. The original idea to use Hilbert's metric in the analysis of cone mappings is due to Garrett Birkhoff [3] and H. Samelson [26], who used it to analyze eigenvalue problems for linear operators that leave a closed cone in a Banach space invariant.

Interesting nonlinear examples arise in optimal control and game theory [7], [25], matrix scaling problems [23], and the analysis of diffusions on fractals [18], [21]. In these applications it is often important to understand the iterative behavior of  $g: X_\phi \rightarrow X_\phi$  in (1.1).

The goal of this paper is to analyze the iterates of mappings  $g: X \rightarrow X$  which are nonexpansive with respect to Hilbert's metric, for strictly convex domains  $X$ , meaning that  $\partial X$  does not contain any straight-line segments, and for  $X$  an open  $n$ -simplex. It is important to distinguish two cases:  $g$  has a fixed point in  $X$ , and  $g$  does not have a fixed point in  $X$ . In the second case it follows from a result by A. Calka [6] that the limit points of each orbit of  $g$  are contained

in the boundary of  $X$ . In fact, Nussbaum [24] and Karlsson have conjectured independently that, in that case, there exists  $\Lambda \subseteq \partial X$  convex such that the limit points of all orbits of  $g$  are contained in  $\Lambda$ . If  $X$  is a strictly convex, the conjecture is known to be true [2] and  $\Lambda$  reduces to a single point. Partial results for general domains were obtained in [2], [12], [13], [17], [24].

In this paper we shall restrict ourselves to the first case and assume that  $g$  has a fixed point in  $X$ . We prove the following result.

**THEOREM 1.1.** *Suppose that  $(X, d_X)$  is a strictly convex Hilbert's metric space and there exists no 2-dimensional plane  $P$  such that  $X \cap P$  is the interior of an ellipse. If  $g: X \rightarrow X$  is nonexpansive under Hilbert's metric and  $g$  has a fixed point, then every orbit of  $g$  converges to a periodic orbit. In fact, there exists an integer  $q \geq 1$  such that  $(g^{qk}(x))_k$  converges for all  $x \in X$ .*

In other words, each orbit of a nonexpansive mapping converges to a periodic orbit if the domain does not contain a hyperbolic plane and the mapping has a fixed point.

If the domain  $X$  is an open polytope, i.e. the intersection of finitely many open half-spaces, then it is known [22], [27] that there is convergence to periodic orbits for nonexpansive mappings  $g: X \rightarrow X$  with a fixed point in  $X$ . Moreover, there exists an a priori upper bound for the possible periods in terms of the number of facets of  $X$ . The current best estimate, obtained in [16], is

$$\max_{k=1, \dots, m} 2^k \binom{m}{k},$$

where  $m = N(N-1)/2$  and  $N$  is the number of facets of  $X$ . This upper bound is believed to be far from optimal. Finding a sharp upper bound appears to be a hard combinatorial geometric problem, even when  $X$  is an open  $n$ -simplex. For the 2-simplex, however, we prove the following result.

**THEOREM 1.2.** *If  $X$  is an open 2-simplex, then the optimal upper bound for the periods of periodic points of nonexpansive mappings on  $(X, d_X)$  is 6.*

## 2. Preliminaries

Let  $(Y, d)$  be a complete metric space, and  $g: Y \rightarrow Y$  be a continuous map. The orbit of  $y$  under  $g$  is denoted by  $\mathcal{O}(y) = \{g^k(y): k = 0, 1, \dots\}$ . A point  $y \in Y$  is called a *periodic point* if  $g^p(y) = y$  for some integer  $p \geq 1$ , and the smallest such  $p \geq 1$  is called the *period* of  $y$ . For  $y \in Y$  the  $\omega$ -limit set of  $y$  under  $g$  is given by,

$$\omega(y; g) = \left\{ x \in Y: \lim_{i \rightarrow \infty} g^{k_i}(y) = x \text{ for some subsequence } k_i \rightarrow \infty \right\}.$$

Furthermore we write  $\Omega_g = \bigcup_{y \in Y} \omega(y; g)$  to denote the *attractor* of  $g$ .

Clearly  $\omega(y; g)$  is closed and  $g(\omega(y; g)) \subseteq \omega(y; g)$ . It is not hard to show that if  $g: Y \rightarrow Y$  is continuous,  $\mathcal{O}(y)$  is pre-compact, and  $|\omega(y; g)| = q$ , then  $(g^{qk}(y))_k$  converges to a periodic point of  $g$  with period  $q$ . Moreover, if  $g: Y \rightarrow Y$  is nonexpansive,  $g$  has a fixed point in  $Y$ , and the orbit of each point in  $Y$  is pre-compact, then each  $\omega(y; g)$  is a non-empty compact set, and  $g(\omega(y; g)) = \omega(y; g)$ . Furthermore it was shown in [8] that  $\omega(x; g) = \omega(y; g)$  for all  $x \in \omega(y; g)$ , and the restriction of  $g$  to  $\omega(y; g)$  is an isometry, see [10].

A metric space  $(Y, d)$  is called *proper* if every closed ball is compact. Hilbert's metric spaces are separable and proper, since their topology coincides with the norm topology, see [22]. As the iterates of a nonexpansive mapping form an equicontinuous family, one can use an Arzelà–Ascoli type argument to prove the following assertion, see [4, p. 9] for details.

**LEMMA 2.1.** *If  $(Y, d)$  is a separable proper metric space and  $g: Y \rightarrow Y$  is a nonexpansive mapping with a fixed point in  $Y$ , then every subsequence of  $(g^k)_k$  has a further subsequence which converges uniformly on compact subsets of  $Y$ .*

Lemma 2.1 can be used to prove the following proposition concerning the existence of a nonexpansive retraction on  $\Omega_g$ . The argument is a straightforward adaptation of [15, Proposition 2.1].

**PROPOSITION 2.2.** *If  $(Y, d)$  is separable proper metric space and  $g: Y \rightarrow Y$  is a nonexpansive mapping with a fixed point on  $Y$ , then there exists a nonexpansive retraction  $r: Y \rightarrow Y$  onto  $\Omega_g$  and the restriction of  $g$  to  $\Omega_g$  is an isometry.*

### 3. Strictly convex domains

Recall that a path  $\gamma: [r, s] \rightarrow (X, d_X)$  is called a *geodesic* if

$$d_X(\gamma(t_1), \gamma(t_2)) = |t_1 - t_2| \quad \text{for all } t_1, t_2 \in [r, s].$$

A *geodesic segment* is the image of a geodesic in  $(X, d_X)$ . It is a well-known fact that if  $X$  is strictly convex, then  $(X, d_X)$  is uniquely geodesic [4, p. 106]; so, the only geodesic segments are the straight-line segments. For  $x, y \in (X, d_X)$  we write  $[x, y]$  to denote the straight-line segment connecting  $x$  and  $y$ , and  $\ell_{x,y}$  to denote the straight-line through  $x$  and  $y$ .

**LEMMA 3.1.** *If  $(X, d_X)$  is a strictly convex Hilbert's metric space and  $g: X \rightarrow X$  is a nonexpansive mapping with a fixed point in  $X$ , then  $\Omega_g$  is convex.*

**PROOF.** By Proposition 2.2 there exists a nonexpansive retraction  $r: X \rightarrow X$  onto  $\Omega_g$ . Thus,  $r(r(x)) = r(x)$  for all  $x \in X$ .

Now let  $x, y \in \Omega_g$  and put  $s = d_X(x, y)$ . Let  $\gamma: [0, s] \rightarrow X$  be the unique geodesic path connecting  $[x, y]$ , so the image of  $\gamma$  is the straight-line segment

between  $x$  and  $y$ . Now let  $u = \gamma(t)$ , so  $d_X(x, u) = t$  and  $d_X(u, y) = s - t$ . As  $r$  is nonexpansive, we get that  $d_X(x, r(u)) \leq t$  and  $d_X(y, r(u)) \leq s - t$ . Now using

$$d_X(x, r(u)) + d_X(r(u), y) \leq d_X(x, y) = s,$$

we find that  $d_X(x, r(u)) = t$  and  $d_X(y, r(u)) = s - t$ . This implies that  $r(u)$  lies on the unique geodesic connecting  $x$  and  $y$ , which is  $[x, y]$ . Thus,  $r(u) = u$  and hence  $u \in \Omega_g$ .  $\square$

It is convenient to embed the domain  $X$  into the affine hyperplane  $H = \{(x, 1) \in \mathbb{R}^{n+1} : x \in \mathbb{R}^n\}$  in  $\mathbb{R}^{n+1}$  by identifying  $X$  with  $\{(x, 1) \in \mathbb{R}^{n+1} : x \in X\}$ . Let  $P(\mathbb{R}^{n+1})$  denote the real  $n$ -dimensional projective space. So,  $P(\mathbb{R}^{n+1})$  is the set of lines through the origin in  $\mathbb{R}^{n+1}$ . Recall that  $P(\mathbb{R}^{n+1})$  can be partitioned into the set of lines intersecting the hyperplane  $H$  and the set of lines parallel to  $H$ . In other words,  $X$  is contained in the open cell of  $P(\mathbb{R}^{n+1})$ . Let  $V' = \text{aff}(\Omega_g)$  denote the affine span of  $\Omega_g$  inside  $H$ , and put  $X' = V' \cap X$ . Let  $V$  be the vector space above  $V'$  in  $\mathbb{R}^{n+1}$ . Thus,  $X'$  lies in the open cell of the real projective space  $P(V) = V' \cup P(V')$ . We write

$$\text{Coll}(X') = \{h \in \text{PGL}(V) : h(X') = X'\}$$

to denote the *collineation group* of  $X'$ .

**PROPOSITION 3.2.** *If  $g : X \rightarrow X$  is a nonexpansive mapping on a strictly convex Hilbert's metric space  $(X, d_X)$ , and  $g$  has a fixed point, then there exists a collineation  $h \in \text{Coll}(X')$  such that  $h$  coincides with  $g$  on  $\Omega_g$ .*

**PROOF.** Let  $k = \dim V' \geq 1$ . (The case  $k = 0$  is trivial.) By Lemma 3.1  $\Omega_g$  is convex, and hence there exist  $x^0, \dots, x^k$  in  $\Omega_g$  such that  $\text{conv}(x^0, \dots, x^k)$  is a  $k$ -simplex in  $\Omega_g$ . Let  $y \in \Omega_g$  be in the relative interior of  $\text{conv}(x^0, \dots, x^k)$ . So,  $y, x^0, \dots, x^k$  forms a projective basis for  $P(V)$ .

As  $g$  is an isometry on  $\Omega_g$  and  $g$  maps  $\Omega_g$  onto itself,  $g|_{\Omega_g}$  and  $g|_{\Omega_g}^{-1}$  map straight-line segments to straight-line segments in  $\Omega_g$ . We now use a simple induction argument to prove the following claim.

*Claim 1.* For  $0 \leq k_0 < \dots < k_s \leq k$  we have that  $z \in \text{conv}(x^{k_0}, \dots, x^{k_s})$  if and only if  $g(z) \in \text{conv}(g(x^{k_0}), \dots, g(x^{k_s}))$ .

If  $s = 1$ , then the assertion is clear as  $g$  maps the straight-line segment  $[x^{k_0}, x^{k_1}]$  onto the straight-line segment  $[g(x^{k_0}), g(x^{k_1})]$ . Now suppose the assertion is true for all integers  $r$  with  $1 \leq r < s \leq k$ . Let  $z \in \text{conv}(x^{k_1}, \dots, x^{k_s})$ . The assertion is clearly true for  $z = x^{k_s}$ . So, suppose that  $z \neq x^{k_s}$  and let  $\ell$  be the straight-line through  $z$  and  $x^{k_s}$ . Note that  $\ell$  intersects  $\text{conv}(x^{k_1}, \dots, x^{k_{s-1}})$  in a point  $u$ . By the induction hypothesis  $g(u) \in \text{conv}(g(x^{k_1}), \dots, g(x^{k_{s-1}}))$ . As  $g$  maps  $[u, x^{k_s}]$  onto  $[g(u), g(x^{k_s})]$ , we conclude that  $g(z)$  is in  $\text{conv}(g(x^{k_0}), \dots,$

$g(x^{k_s}))$ . The opposite implication is obtained by applying the same argument to  $g|_{\Omega_g}^{-1}$ .

Claim 1 implies that  $\text{conv}(g(x^0), \dots, g(x^k))$  is a  $k$ -dimensional simplex in  $\Omega_g$  and  $g(y)$  is in its relative interior. Thus,  $g(y), g(x^0), \dots, g(x^k)$  is a projective basis for  $P(V)$ . Let  $h \in \text{PGL}(V)$  be the unique collineation that coincides with  $g$  on  $y, x^0, \dots, x^k$ . We will show that  $h \in \text{Coll}(X')$  and  $h$  coincides with  $g$  on the whole of  $\Omega_g$ . We need the following claim.

*Claim 2.* If  $g$  coincides with  $h$  on 3 distinct collinear points  $x, w, z \in \Omega_g$ , then  $g$  coincides with  $h$  on the straight-line segment  $\ell_{x,z} \cap \Omega_g$ .

To prove the claim let  $a, b \in \partial X'$  be the points of intersection of the straight-line through  $x$  and  $z$  such that  $x$  is between  $a$  and  $z$ , and  $z$  is between  $b$  and  $x$ . Likewise let  $a', b' \in \partial X'$  be the points of intersection of the straight-line through  $g(x)$  and  $g(z)$  such that  $g(x)$  is between  $a'$  and  $g(z)$ , and  $g(z)$  is between  $b'$  and  $g(x)$ . There exists a collineation  $f$  on the projective line containing  $\ell_{x,z}$  that maps  $a$  to  $a'$ ,  $b$  to  $b'$ , and  $x$  to  $g(x)$ . We show that  $f$  coincides with  $g$  on  $\ell_{x,z} \cap \Omega_g$ . For  $u \in \ell_{x,z} \cap \Omega_g$ , with  $u$  between  $x$  and  $b$ ,

$$[a', g(x), g(u), b'] = [a, x, u, b] = [a', f(x), f(u), b'].$$

As  $f(x) = g(x)$ , this equality uniquely determines  $g(u)$ , so that  $g(u) = f(u)$ . The case where  $u$  is between  $z$  and  $a$  is similar. Now note that, as  $x, w$  and  $z$  form a projective basis for the projective line containing  $\ell_{x,z}$ , and  $h$  and  $f$  coincides on these 3 points,  $f$  and  $h$  are identical on  $\ell_{x,z}$ . This implies that  $g$  and  $h$  are identical on  $\ell_{x,z} \cap \Omega_g$ .

Note that for each  $0 \leq l \leq k$  there exists  $y^l$  in the relative interior of  $\text{conv}(\{x^1, \dots, x^k\} \setminus \{x^l\})$  such that  $h(y^l) = g(y^l)$ . Simply let  $y^l$  be the point of intersection of  $\ell_{x^l, y}$  and  $\text{conv}(\{x^1, \dots, x^k\} \setminus \{x^l\})$ . It follows from Claim 1 that  $g(y^l)$  is in the relative interior of  $\text{conv}(\{g(x^0), \dots, g(x^k)\} \setminus \{g(x^l)\})$ . As  $g$  maps straight-line segments to straight-line segments.  $g(y^l)$  is the unique point of intersection of  $\ell_{g(x^l), g(y)}$  and  $\text{conv}(\{g(x^0), \dots, g(x^k)\} \setminus \{g(x^l)\})$ . Thus,  $h(y^l) = g(y^l)$ . Repeating this argument shows that for each  $0 \leq k_0 < k_1 < \dots < k_s \leq k$  there exists  $w$  in the relative interior of  $\text{conv}(x^{k_0}, \dots, x^{k_s})$  such that  $h(w) = g(w)$ .

We shall now show by induction on  $s \geq 1$  that  $g$  and  $h$  coincide on  $\text{conv}(x^{k_0}, \dots, x^{k_s})$ . The induction basis is true by Claim 2. Suppose the assertion is true for all  $m < s$ . Let  $v$  and  $w$  be points in the relative interior of  $\text{conv}(x^{k_0}, \dots, x^{k_s})$  with  $v \neq w$  and  $h(w) = g(w)$ . Then the straight-line  $\ell_{w,v}$  intersects the relative boundary of  $\text{conv}(x^{k_0}, \dots, x^{k_s})$  in two distinct points  $p$  and  $q$ . By the induction hypothesis  $g$  and  $h$  coincide on  $p$  and  $q$ . As  $h(w) = g(w)$ , we can apply Claim 2 to deduce that  $g(v) = h(v)$ .

To see that  $g$  coincides with  $h$  on the whole of  $\Omega_g$  we remark that for  $v \in \Omega_g$  the straight-line  $\ell_{y,v}$  contains at least 3 points of  $\text{conv}(x^0, \dots, x^k)$ , as  $y$  is in the relative interior of  $\text{conv}(x^0, \dots, x^k)$ . Thus, by Claim 2  $g$  coincides with  $h$  on  $\Omega_g$ .

It remains to show that  $h(X') = X'$ . Let  $z_1 \neq z_2$  in the relative interior of  $\text{conv}(x^0, \dots, x^k)$ , and let  $a, b \in \partial X' \cap \ell_{z_1, z_2}$  such that  $z_1$  is between  $z_2$  and  $a$ , and  $z_2$  is between  $z_1$  and  $b$ . Then  $[a, z_1, z_2, b] = [a', g(z_1), g(z_2), b']$ , where  $a', b' \in \partial X' \cap \ell_{g(z_1), g(z_2)}$ , and  $g(z_1)$  is between  $a'$  and  $g(z_2)$ . There exists a collineation  $f$  that maps  $a$  to  $a'$ ,  $b$  to  $b'$ , and  $z_1$  to  $g(z_1)$ . As in the proof of Claim 2  $g$  coincides with  $f$  on  $[z_1, z_2]$ . This implies that  $h$  coincides with  $f$  on  $\ell_{z_1, z_2} \cap X'$ , and hence  $h(a) = a'$  and  $h(b) = b'$ , which completes the proof.  $\square$

Ideas similar to the ones behind Lemma 3.1 and Proposition 3.2 were used by Edelstein in [9] to analyze nonexpansive mappings on strictly convex Banach spaces.

A key ingredient in the proof of Theorem 1.1 is a result by Y. I. Lyubich and A. I. Veitsblit [20]. To state it we need to recall a few definitions. Let  $C$  be a closed cone with non-empty interior in a finite dimensional real vector space  $W$ , so  $C$  is convex,  $\lambda C \subseteq C$  for all  $\lambda > 0$ , and  $C \cap (-C) = \{0\}$ . A linear map  $A: W \rightarrow W$  is said to be *positive* if  $A(C) \subseteq C$ . We denote the *automorphism group* of  $C$  by

$$\text{Aut}(C) = \{A \in \text{GL}(W) : A(C) = C\}.$$

A linear subspace  $U$  of  $W$  is called  *$C$ -complemented* if there exists a positive linear projection  $P: W \rightarrow W$  with range  $U$ . Note that if  $U$  is a subspace of  $W$ , then  $U \cap C$  is a closed sub-cone of  $C$ .

A closed cone  $C$  with non-empty interior in  $W$ , where  $\dim W = n + 1$ , is called a *Lorentz cone* if there exists a system of coordinates  $x_1, \dots, x_{n+1}$  such that

$$C = \{(x_1, \dots, x_{n+1}) \in W : x_{n+1}^2 - x_1^2 - \dots - x_n^2 \geq 0 \text{ and } x_{n+1} \geq 0\}.$$

**THEOREM 2.3** (Lyubich–Veitsblit [20]). *If  $C \subseteq W$  is a closed cone with non-empty interior, and  $\text{Aut}(C)$  contains an infinite compact subgroup, then there exists a 3-dimensional  $C$ -complemented subspace  $U$  of  $W$  such that  $K = C \cap U$  is a Lorentz cone in  $U$ .*

We will now prove Theorem 1.1.

**PROOF OF THEOREM 1.1.** As before identify  $X$  with  $\{(x, 1) \in \mathbb{R}^{n+1} : x \in X\}$  and let  $V' = \text{aff}(\Omega_g)$  be the affine span of  $\Omega_g$  in  $\mathbb{R}^{n+1}$ . Put  $X' = V' \cap X$  and let  $V$  be the subspace above  $V'$  in  $\mathbb{R}^{n+1}$ .

Furthermore let  $C_X$  denote the closure of the open cone generated by  $X$  in  $\mathbb{R}^{n+1}$ , so

$$C_X = \{\lambda(x, 1) \in \mathbb{R}^{n+1} : x \in \overline{X} \text{ and } \lambda \geq 0\}.$$

Likewise let  $C_{X'}$  denote the closure of the open cone generated by  $X'$  in  $V$ . Let  $x^* \in \Omega_g$  be a fixed point of  $g$ , and note that  $x^*$  is in the interior of  $C_{X'}$  in  $V$ .

By Proposition 2.2 there exists an  $h \in \text{Coll}(X')$  such that  $h$  coincides with  $g$  on  $\Omega_g$ . Thus, there exists  $A \in \text{Aut}(C_{X'})$  such that  $pA = h$ , where  $p$  denotes the projection. Remark that, as  $h(x^*) = g(x^*) = x^*$ , we have that  $A(x^*) = \sigma x^*$  for some  $\sigma > 0$ . Putting  $B = \frac{1}{\sigma}A$  we see that  $B \in \text{Aut}(C_{X'})$ ,  $pB = h$  and  $B(x^*) = x^*$ .

As  $x^*$  is in the interior of  $C_{X'}$  in  $V$ , we know that  $(B^k)_{k=0}^\infty$  is bounded. Indeed, let  $\|x\| = \inf\{\mu > 0: -\mu x^* \leq_C x \leq_C \mu x^*\}$  be the order unit norm on  $V$ , see [22, p.14]. Now let  $x \in V$  and  $\|x\| = \tau$ . Then  $-\tau x^* \leq_C x \leq_C \tau x^*$ , so that

$$-\tau x^* = -\tau B(x^*) \leq_C B(x) \leq_C \tau B(x^*) = \tau x^*.$$

This implies that  $\|B\| = 1$ , and hence  $(B^k)_{k=0}^\infty$  is bounded. In the same way it can be shown that  $\|B^{-1}\| = 1$ , and hence the closure of the group generated by  $(B^k)_{k=0}^\infty$  is a compact subgroup of  $\text{Aut}(C_{X'})$ .

As there exists no 2-dimensional plane  $P$  in  $\mathbb{R}^n$  such that  $P \cap X$  is the interior of an ellipse, there exists no 3-dimensional subspace  $U$  of  $\mathbb{R}^{n+1}$  such that  $U \cap C_{X'}$  is a Lorentz cone. Hence by the Lyubich–Veitsblit Theorem 2.3 we know that there exists an integer  $q \geq 1$  such that  $B^q = I$ .

Now to prove the convergence to periodic orbits, it suffices to show that  $|\omega(x; g)|$  divides  $q$  for each  $x \in X$ , since  $g$  is nonexpansive. So, let  $x \in X$  and  $y \in \omega(y; g)$ . By [8] we know that  $\omega(x; g) = \omega(y; g)$ . As  $y \in \Omega_g$ ,  $g^q(y) = h^q(y) = pB^q(y) = y$ . Therefore  $y$  is a periodic point of  $g$  whose period divides  $q$ . Thus,  $|\omega(x; g)| = |\omega(y; g)|$  divides  $q$ , and we are done.  $\square$

The proof of Theorem 1.1 does not use the full strength of the Lyubich–Veitsblit Theorem 2.3. In fact, I believe that the hypothesis in Theorem 1.1 can be weakened to the assumption: there exists no 3-dimensional  $C_X$ -complemented subspace of  $\mathbb{R}^{n+1}$  such that  $C_X \cap U$  is a Lorentz cone. For instance, if  $C_X = \{(x_1, \dots, x_4) \in \mathbb{R}^4 : x_4^4 - x_1^4 - x_2^4 - x_3^4 \geq 0 \text{ and } x_4 \geq 0\}$ , which corresponds to  $X$  being the interior of the unit ball in  $\mathbb{R}^3$  with the  $\ell_4$ -norm, then it is known [19] that there exists a 3-dimensional subspace  $U$  such that  $C_X \cap U$  is a Lorentz cone, but  $U$  is not  $C_X$ -complemented, see [20, Theorem 3]. A more daring speculation would be to conjecture that there is convergence to periodic orbits for general Hilbert's metric spaces not containing a hyperbolic plane.

It must be noted that there exists an analogous result for finite dimensional strictly convex normed spaces, see [15]. In that case one assumes that there exists no 1-complemented Euclidean plane to ensure convergence to periodic orbits.



### 4. The simplex

It is known [22] that if  $X$  is an open  $n$ -simplex, then  $(X, d_X)$  is isometric to a normed space  $(\mathbb{R}^n, \|\cdot\|_H)$ , where  $\|\cdot\|_H$  has a polyhedral unit ball. As  $(X, d_X)$  and  $(Y, d_Y)$  are isometric if  $X$  and  $Y$  are open  $n$ -simplices, we can restrict ourselves to analyzing Hilbert's metric on the *standard  $n$ -simplex*,

$$\Delta_n = \left\{ (x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : \sum_i x_i = 1 \text{ and } x_i > 0 \text{ for } i = 1, \dots, n+1 \right\}.$$

In that case it is easy to describe the isometry. Let  $\mathbb{R}^{n+1}$  be equipped with an equivalence relation  $\sim$  given by,  $x \sim y$  if  $x = y + \lambda(1, \dots, 1)$  for some  $\lambda \in \mathbb{R}$ . Then  $\mathbb{R}^{n+1}/\sim$  is an  $n$ -dimensional vector space that can be endowed with the *variation norm*,

$$\|x\|_{\text{var}} = \max_{1 \leq i \leq n+1} x_i - \min_{1 \leq j \leq n+1} x_j \quad \text{for } x \in \mathbb{R}^{n+1}/\sim.$$

There exists an isometry of  $(\Delta_n, d_{\Delta_n})$  onto  $(\mathbb{R}^{n+1}/\sim, \|\cdot\|_{\text{var}})$  which is given by,

$$\text{Log}(x) = (\log x_1, \dots, \log x_{n+1}) \quad \text{for } x \in \Delta_n.$$

By taking the representative  $x \in \mathbb{R}^{n+1}$  with  $x_{n+1} = 0$  in the equivalence class of  $x$  in  $\mathbb{R}^{n+1}/\sim$ , and projecting out the  $(n+1)$ -th coordinate, we see that  $(\Delta_n, d_{\Delta_n})$  is isometric to  $(\mathbb{R}^n, \|\cdot\|_H)$ , where

$$\|z\|_H = \left( 0 \vee \max_{1 \leq i \leq n} z_i \right) - \left( 0 \wedge \min_{1 \leq j \leq n} z_j \right) \quad \text{for } z \in \mathbb{R}^n.$$

Here  $a \vee b = \max\{a, b\}$  and  $a \wedge b = \min\{a, b\}$ . In dimension  $n = 2$  the unit ball is a hexagon, as shown in Figure 2. For  $n = 3$  the unit ball is a rhombic-dodecahedron.




FIGURE 2. The unit ball of  $\|\cdot\|_H$

It is clear that there exists an isometry  $A$  on  $(\mathbb{R}^n, \|\cdot\|_H)$  with period 6. Simply take

$$A = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}.$$

Thus, on  $(\Delta_2, d_{\Delta_2})$  there exists a nonexpansive mapping that has a period 6 point. In this section it is shown that 6 is the maximal possible period. The proof uses similar methods as the ones developed in [14].

Let  $(\mathbb{R}^n, \|\cdot\|)$  be a polyhedral normed space, i.e. its unit ball is a polyhedron. A sequence  $x^1, \dots, x^k \in \mathbb{R}^n$  is called an *additive chain* if

$$\|x^1 - x^k\| = \sum_{i=1}^{k-1} \|x^i - x^{i+1}\|.$$

As  $\|\cdot\|$  is a polyhedral norm,  $x^1, x^2, \dots, x^k$  need not lie on a straight-line in order to be an additive chain. For  $x, y \in \mathbb{R}^n$  define

$$W(x, y) = \{z \in \mathbb{R}^n : x, y, z \text{ is an additive chain}\}$$

and denote its interior by  $W^\circ(x, y)$ . Given a polyhedral norm on  $\mathbb{R}^n$  whose unit ball has  $m$  facets, i.e.  $m$  faces of dimension  $n-1$ , there exist  $m$  linear functionals  $\phi_1, \dots, \phi_m$  such that

$$\|x\| = \max_{i=1, \dots, m} \langle \phi_i, x \rangle \quad \text{for all } x \in \mathbb{R}^n.$$

Let  $I(x, y) = \{i : \|x - y\| = \phi_i(x - y)\}$ .

**LEMMA 4.1.** *If  $x^1, \dots, x^k$  is a sequence in a polyhedral normed vector space  $(\mathbb{R}^n, \|\cdot\|)$  and  $\|x^1 - x^k\| = \phi_i(x^1 - x^k)$ , then  $x^1, \dots, x^k$  is an additive chain if and only if  $\|x^j - x^{j+1}\| = \phi_i(x^j - x^{j+1})$  for all  $j = 1, \dots, k-1$ .*

**PROOF.** Clearly

$$\|x^1 - x^k\| = \sum_{j=1}^{k-1} \|x^j - x^{j+1}\| \geq \sum_{j=1}^{k-1} \phi_i(x^j - x^{j+1}) = \phi_i(x^1 - x^k) = \|x^1 - x^k\|.$$

Thus,  $\phi_i(x^j - x^{j+1}) = \|x^j - x^{j+1}\|$  for all  $j = 1, \dots, k-1$ . Conversely,

$$\|x^1 - x^k\| \leq \sum_{j=1}^{k-1} \|x^j - x^{j+1}\| = \sum_{j=1}^{k-1} \phi_i(x^j - x^{j+1}) = \phi_i(x^1 - x^k) \leq \|x^1 - x^k\|. \quad \square$$

It follows from Lemma 4.1 that  $x^1, x^2, \dots, x^k$  is an additive chain if and only if  $\bigcap_{j=1}^{k-1} I(x^j, x^{j+1}) \neq \emptyset$ .

LEMMA 4.2. *For  $x \neq y$  in a polyhedral normed vector  $(\mathbb{R}^n, \|\cdot\|)$  we have that*

$$W^\circ(x, y) = \{z \in W(x, y) : I(y, z) \subseteq I(x, y)\}.$$

PROOF. Suppose that  $x, y, z$  is an additive chain and  $I(y, z) \not\subseteq I(x, y)$ . Let  $i \in I(y, z) \setminus I(x, y)$ . For  $\varepsilon > 0$  there exists  $z' \in \mathbb{R}^n$  such that  $I(y, z') = \{i\}$  and  $\|z' - z\| \leq \varepsilon$ . Note that  $x, y, z'$  is not an additive chain, as  $I(x, y) \cap I(y, z') = \emptyset$ . Hence  $z \in \partial W(x, y)$ , which shows that

$$W^\circ(x, y) \subseteq \{z \in W(x, y) : I(y, z) \subseteq I(x, y)\}.$$

On the other hand, given an additive chain  $x, y, z$ , we let

$$\varepsilon = \min_{i,j} \phi_i(y - z) - \phi_j(y - z) > 0,$$

where the minimum is taken over all  $i \in I(y, z)$  and  $j \notin I(y, z)$ .

For each  $z' \in \mathbb{R}^n$  with  $\|z - z'\| < \varepsilon/2$ , we have that  $I(y, z') \subseteq I(y, z)$ . Indeed, if  $j \notin I(y, z)$  and  $i \in I(y, z)$ , then

$$\begin{aligned} \phi_j(y - z') &= \phi_j(y - z) + \phi_j(z - z') \leq \phi_i(y - z) + \phi_j(z - z') - \varepsilon < \phi_i(y - z) - \varepsilon/2 \\ &\leq \phi_i(y - z') + \phi_i(z' - z) - \varepsilon/2 < \phi_i(y - z'), \end{aligned}$$

and hence  $j \notin I(y, z')$ .

As  $I(y, z) \subseteq I(x, y)$ , we know that  $I(y, z') \subseteq I(x, y)$ . This implies that  $I(y, z') \cap I(x, y)$  is non-empty, and hence  $x, y, z'$  is an additive chain.  $\square$

Recall that if  $\mathcal{O} = \{\xi, g(\xi), \dots, g^{p-1}(\xi)\}$  is a periodic orbit of a nonexpansive mapping  $g$  on a metric space  $(Y, d)$ , then the iterates  $\Gamma = \{g^k : k = 0, \dots, p-1\}$  form a cyclic group of isometries on  $\mathcal{O}$  that acts transitively on  $\mathcal{O}$ , i.e. for each  $x, y \in \mathcal{O}$  there exists  $g^k \in \Gamma$  such that  $g^k(x) = y$ . The following lemma generalizes [14, Lemma 2.2].

LEMMA 4.3. *If  $S$  is a compact set in a polyhedral normed space  $(\mathbb{R}^n, \|\cdot\|)$  and  $S$  has a transitive commutative group of isometries, then*

$$W^\circ(x, y) \cap S = \emptyset \quad \text{for all } x \neq y \in S.$$

PROOF. Let  $x, y, z \in S$  be such that  $x \neq y$  and  $z \in W^\circ(x, y)$ . Put  $\varepsilon = \min\{\|x - y\|, \|y - z\|\} > 0$ . Denote the collection of all additive chains in  $S$  starting with  $x, y, z$  and which are such that the distance between consecutive points in the chain is at least  $\varepsilon$  by  $\mathcal{F}$ .

Note that, since  $S$  is compact, there exists an upper bound on the length of the additive chains in  $\mathcal{F}$ . Let  $x^1 = x, x^2 = y, x^3 = z, \dots, x^r$  be an additive chain in  $\mathcal{F}$  of maximal length.

For  $1 \leq k, l \leq r$  let  $g_{k,l}$  be an isometry on  $S$  in the commutative group that acts transitively on  $S$  such that  $g_{k,l}(x^k) = x^l$ . Denote  $x^{r+1} = g_{1,2}(x^r)$ . We now

show that  $x^2, x^3, \dots, x^{r+1}$  is also an additive chain in  $S$  in which the distance between consecutive points is at least  $\varepsilon$ . Indeed, as the group is commutative,

$$\|x^r - x^{r+1}\| = \|g_{1,r}(x^1) - g_{1,r}(g_{1,2}(x^1))\| = \|x^1 - x^2\| \geq \varepsilon.$$

This equality implies,

$$\|x^2 - x^{r+1}\| = \|g_{1,2}(x^1) - g_{1,2}(x^r)\| = \sum_{j=1}^{r-1} \|x^j - x^{j+1}\| = \sum_{j=2}^r \|x^j - x^{j+1}\|,$$

and hence  $\bigcap_{j=2}^r I(x^j, x^{j+1}) \neq \emptyset$ . As  $z \in W^\circ(x, y)$ , we know that  $I(x^2, x^3) \subseteq I(x^1, x^2)$ , and hence  $\bigcap_{j=1}^r I(x^j, x^{j+1}) \neq \emptyset$ . But this implies that  $x^1, \dots, x^{r+1}$  is an additive chain in  $\mathcal{F}$ , which contradicts the maximality of  $r$ .  $\square$

We can now prove Theorem 1.2.

PROOF OF THEOREM 1.2. If  $g$  is a nonexpansive mapping on  $(\mathbb{R}^2, \|\cdot\|_H)$  with a periodic orbit  $\mathcal{O}$  and fixed point  $x^*$ , then  $\mathcal{O}$  is contained in the boundary of the ball  $B(x^*) = \{x \in \mathbb{R}^2 : \|x - x^*\|_H \leq R\}$  for some  $R \geq 0$ . Without loss of generality we may assume that  $x^* = 0$ .

Partition  $\partial B(x^*)$  as follows:

$$\begin{aligned} A_1 &= \{x \in \mathbb{R}^2 : x_1 = R \text{ and } 0 \leq x_2 < R\}, \\ A_2 &= \{x \in \mathbb{R}^2 : x_2 = R \text{ and } 0 < x_1 \leq R\}, \\ A_3 &= \{x \in \mathbb{R}^2 : x_2 - x_1 = R \text{ and } 0 < x_2 < R\}, \\ A_4 &= \{x \in \mathbb{R}^2 : x_1 = -R \text{ and } -R < x_2 \leq 0\}, \\ A_5 &= \{x \in \mathbb{R}^2 : x_2 = -R \text{ and } -R \leq x_1 < 0\}, \\ A_6 &= \{x \in \mathbb{R}^2 : x_1 - x_2 = R \text{ and } -R \leq x_2 < 0\}. \end{aligned}$$

Each  $A_i$  corresponds to a facet of  $\partial B(x^*)$  with one of its vertices removed.

By Lemma 4.3 each  $A_i$  can contain at most two points of  $\mathcal{O}$ , see also Figure 2. Moreover, if  $A_i$  contains 2 points of  $\mathcal{O}$ , then  $A_{i+1} \cap \mathcal{O}$  is empty. (Here we are counting modulo 6.) Indeed, if  $\phi_i$  denotes the facet defining functional of  $\partial B(x^*)$  corresponding to  $A_i$ , then it is easy to verify that if  $x \neq y$  in  $A_i$ , then  $I(x, y) = \{i+1, i+2\}$ , so that  $W^\circ(x, y) \supseteq A_{i+1}$ . Thus,  $A_{i+1} \cap \mathcal{O}$  is empty, if  $x \neq y$  in  $A_i \cap \mathcal{O}$ , and hence  $\mathcal{O}$  has at most 6 points.  $\square$

The example of a period 6 orbit of a nonexpansive mapping on  $\Delta_2$  can be generalized to  $\Delta_n$ . To do this it is convenient to work in  $(\mathbb{R}^{n+1}/\sim, \|\cdot\|_{\text{var}})$ . Let  $\mathcal{X}$  be the subset of all points  $x \in \mathbb{R}^{n+1}/\sim$  such that  $x_i = 1$  or  $x_i = 0$  for each  $i$ . Note that  $(0, \dots, 0) = (1, \dots, 1)$  and that for each  $x \in \mathcal{X}$  we have  $-x = (1, \dots, 1) - x \in \mathcal{X}$ . If  $\mathcal{S} \subseteq \mathcal{X}$  is such that there exists no  $x \neq y$  in  $\mathcal{S}$  with  $x \leq y$ , where the inequality holds coordinate-wise, then  $\|x - y\|_{\text{var}} = 2$  for all  $x \neq y$  in  $\mathcal{S}$ . For such sets  $\mathcal{S}$  it is known [16, p. 867] that there exists

a nonexpansive mapping  $g_S$  on  $\mathbb{R}^{n+1}/\sim$  which has  $S$  as a periodic orbit. Now suppose that  $n$  is even and let  $\mathcal{S}_n \subseteq \mathcal{X}$  be the set of points with exactly  $n/2$  coordinates equal to 1. Then  $-\mathcal{S}_n$  is the set of points with exactly  $n/2 + 1$  coordinates equal to 1 in  $\mathcal{X}$ . Clearly the reflection in the origin,  $R(x) = -x$ , is an isometry on  $\mathbb{R}^{n+1}/\sim$ . Now if  $|\mathcal{S}_n| = \binom{n+1}{n/2}$  is odd, the nonexpansive mapping  $R \circ g_{\mathcal{S}_n}$  has a periodic orbit of length  $2\binom{n+1}{n/2}$ . For  $n = 2$ , this gives a period 6 orbit  $\Delta_2$ . On the other hand, if  $\binom{n+1}{n/2}$  is even, we can drop one point of  $\mathcal{S}_n$  and obtain a periodic orbit of length  $2\binom{n+1}{n/2} - 2$ . A similar construction exists for  $n$  odd. This shows, for general  $n$ , that there exists a nonexpansive mapping on  $(\Delta_n, d_{\Delta_n})$  that has a periodic point with period

$$2\binom{n+1}{\lfloor n/2 \rfloor} - 2\delta_n,$$

where  $\delta_n = 1$  if  $\binom{n+1}{\lfloor n/2 \rfloor}$  is even and  $\delta_n = 0$  otherwise. It seems unlikely that this lower bound is optimal, but no better examples are known.

## REFERENCES

- [1] J. C. ÁLVAREZ PAIVA, *Hilbert's fourth problem in two dimensions*, MASS selecta: teaching and learning advanced undergraduate mathematics (Svetlana Katok et al., eds.), Amer. Math. Soc., Providence, RI, 2003, pp. 165–183.
- [2] A. F. BEARDON, *The dynamics of contractions*, Ergodic Theory Dynam. Systems **17** (1997), 1257–1266.
- [3] G. BIRKHOFF, *Extensions of Jentzsch's theorems*, Trans. Amer. Math. Soc. **85** (1957), 219–277.
- [4] H. BUSEMANN, *The Geometry of Geodesics Pure and Applied Mathematics*, vol. 6, Academic Press, New York, 1955.
- [5] P. BUSHELL, *Hilbert's metric and positive contraction mappings in Banach space*, Arch. Rational Mech. Anal. **52** (1973), 330–338.
- [6] A. CALKA, *On conditions under which isometries have bounded orbits*, Colloq. Math. **48** (1984), 219–227.
- [7] J. COCHET-TERRASSON AND S. GAUBERT, *A policy iteration algorithm for zero-sum stochastic games with mean payoff*, C. R. Math. Acad. Sci. Paris **343** (2006), 377–382.
- [8] C. M. DAFERMOS AND M. SLEMROD, *Asymptotic behavior of nonlinear contraction semigroups*, J. Funct. Anal. **13** (1973), 97–106.
- [9] M. EDELSTEIN, *On non-expansive mappings of Banach spaces*, Proc. Cambridge Philos. Soc. **60** (1964), 439–447.
- [10] H. FREUDENTHAL AND W. HUREWICZ, *Dehnungen, verkürzungen und isometrien*, Fundamenta Math. **26** (1936), 120–122.
- [11] D. HILBERT, *Über die gerade linie als kürzeste verbindung zweier punkte*, Math. Ann. **46** (1895), 91–96.
- [12] A. KARLSSON, *Non-expanding maps and Busemann functions*, Ergodic Theory Dynam. Systems **21** (2001), 1447–1457.
- [13] A. KARLSSON AND G. A. NOSKOV, *The Hilbert metric and Gromov hyperbolicity*, Enseign. Math. **48** (2002), 73–89.

- [14] B. LEMMENS, R. D. NUSSBAUM AND S. M. VERDUYN LUNEL, *Lower and upper bounds for  $\omega$ -limit sets of nonexpansive maps*, Indag. Math. New Ser. **12** (2001), 191–211.
- [15] B. LEMMENS AND O. VAN GAANS, *Dynamics of non-expansive maps on strictly convex Banach spaces*, Israel J. Math. **171** (2009), 425–442.
- [16] B. LEMMENS AND M. SCHEUTZOW, *On the dynamics of sup-norm nonexpansive maps*, Ergodic Theory Dynam. Syst. **25** (2005), 861–871.
- [17] B. LINS, *A Denjoy–Wolff theorem for Hilbert metric nonexpansive maps on polyhedral domains*, Math. Proc. Cambridge Philos. Soc. **143** (2007), 157–164.
- [18] B. LINS AND R. NUSSBAUM, *Denjoy–Wolff theorems, Hilbert metric nonexpansive maps and reproduction-decimation operators*, J. Funct. Anal. **254** (2008), 2365–2386.
- [19] Y. I. LYUBICH AND L. N. VASERSTEIN, *Isometric embeddings between classical Banach spaces, cubature formulas, and spherical designs*, Geom. Dedicata **47** (1993), 327–362.
- [20] YU. I. LYUBICH AND A. I. VEITSBLIT, *Boundary spectrum of nonnegative operators*, Siberian Math. J. **26** (1985), 798–802.
- [21] V. METZ, *Hilbert’s projective metric on cones of Dirichlet forms*, J. Funct. Anal. **127** (1995), 438–455.
- [22] R. D. NUSSBAUM, *Hilbert’s projective metric and iterated nonlinear maps*, Mem. Amer. Math. Soc. **75** (1988), 1–137.
- [22] ———, *Omega limit sets of nonexpansive maps: finiteness and cardinality estimates*, Differential Integral Equations **3** (1990), 523–540.
- [23] ———, *Entropy minimization, Hilbert’s projective metric, and scaling integral kernels*, J. Funct. Anal. **115** (1993), 45–99.
- [24] ———, *Fixed point theorems and Denjoy–Wolff theorems for Hilbert’s projective metric in infinite dimensions*, Topol. Methods Nonlinear Anal. **29** (2007), 199–249.
- [25] D. ROSENBERG AND S. SORIN, *An operator approach to zero-sum repeated games*, Israel J. Math. **121** (2001), 221–246.
- [26] H. SAMELSON, *On the Perron–Frobenius theorem*, Michigan Math. J. **4** (1957), 57–59.
- [27] D. WELLER, *Hilbert’s metric, part metric and selfmappings of a cone*, Ph.D. dissertation (1987), Univ. of Bremen, Germany, December.

*Manuscript received August 4, 2010*

BAS LEMMENS  
 SMSAS  
 University of Kent  
 Canterbury, CT2 7NF, UNITED KINGDOM  
*E-mail address:* B.Lemmens@kent.ac.uk